

Literatur zum Beitrag

Moritz L. Schmidbauer, Andreas Maskos, Stefan Kunst, Konstantinos Dimitriadis

**Sedierung in der Neurointensivmedizin –
Indikationen, therapeutisches
Monitoring und Sedarierungsauslass**

1. S3 Leitlinie Analgesie, Sedierung und Delirmanagement in der Intensivmedizin. https://www.awmf.org/uploads/tx_szleitlinien/001-012I_S3_Analgesie-Sedierung-Delirmanagement-in-der-Intensivmedizin-DAS_2021-08.pdf.
2. Devlin JW, Skrobik Y, Gélinas C et al.: Clinical Practice Guidelines for the Prevention and Management of Pain, Agitation/Sedation, Delirium, Immobility, and Sleep Disruption in Adult Patients in the ICU. *Crit Care Med* 2018; 46(9): e825–e873. doi:10.1097/CCM.0000000000003299
3. Reade MC, Finfer S: Sedation and Delirium in the Intensive Care Unit. *N Engl J Med* 2014; 370(5): 444–54. doi:10.1056/nejmra1208705
4. Girard TD, Kress JP, Fuchs BD et al.: Efficacy and safety of a paired sedation and ventilator weaning protocol for mechanically ventilated patients in intensive care (Awakening and Breathing Controlled trial): a randomised controlled trial. *Lancet* 2008; 371(9607): 126–34. doi:10.1016/S0140-6736(08)60105-1
5. Kress JP, Pohlman AS, O'Connor MF, Hall JB: Daily interruption of sedative infusions in critically ill patients undergoing mechanical ventilation. *N Engl J Med* 2000; 342(20): 1471–7. doi:10.1056/NEJM200005183422002
6. Citerio G, Cormio M: Sedation in neurointensive care: Advances in understanding and practice. *Curr Opin Crit Care* 2003; 9(2): 120–6. doi:10.1097/00075198-200304000-00007
7. Oddo M, Crippa IA, Mehta S et al.: Optimizing sedation in patients with acute brain injury. *Crit Care* 2016; 20(1). doi:10.1186/s13054-016-1294-5
8. Opdenakker O, Vanstraelen A, De Sloovere V, Meyfroidt G: Sedatives in neurocritical care: an update on pharmacological agents and modes of sedation. *Curr Opin Crit Care*. 2019; 25(2): 97–104. doi:10.1097/MCC.0000000000000592
9. Gwinnutt CL, Saha B: Cerebral blood flow and intracranial pressure. *Anaesth Intensive Care Med* 2005; 6(5): 153–6. doi:10.1383/anes.6.5.153.65040
10. Partington T, Farmery A: Intracranial pressure and cerebral blood flow. *Anaesth Intensive Care Med*. 2014; 15(4): 189–94. doi:10.1016/j.mpaic.2014.02.002
11. Carney N, Totten AM, O'Reilly C et al.: Guidelines for the Management of Severe Traumatic Brain Injury, Fourth Edition. *Neurosurgery*. 2017; 80(1): 6–15. doi:10.1227/NEU.0000000000001432
12. Güiza F, Depreitere B, Piper I et al.: Visualizing the pressure and time burden of intracranial hypertension in adult and paediatric traumatic brain injury. *Intensive Care Med*. 2015; 41(6): 1067–76. doi:10.1007/s00134-015-3806-1
13. Hawryluk GWJ, Nielson JL, Huie JR et al.: Analysis of Normal High-Frequency Intracranial Pressure Values and Treatment Threshold in Neurocritical Care Patients: Insights into Normal Values and a Potential Treatment Threshold. *JAMA Neurol* 2020; 77(9): 1150–8. doi:10.1001/jamaneurol.2020.1310
14. Kelly DF, Goodale DB, Williams J et al.: Propofol in the treatment of moderate and severe head injury: a randomized, prospective double-blinded pilot trial. *J Neurosurg* 1999; 90(6): 1042–52. doi:10.3171/jns.1999.90.6.1042
15. Sutter R, Marsch S, Fuhr P, Rüegg S: Mortality and recovery from refractory status epilepticus in the intensive care unit: A 7-year observational study. *Epilepsia* 2013; 54(3): 502–11. doi:10.1111/epi.12064
16. Delaj L, Novy J, Ryvlin P et al.: Refractory and super-refractory status epilepticus in adults: a 9-year cohort study. *Acta Neurol Scand* 2017; 135(1): 92–9. doi:10.1111/ane.12605
17. Rosenow F, Weber J: S2k guidelines: status epilepticus in adulthood: Guidelines of the German Society for Neurology. *Nervenarzt* 2021; 92(10): 1002–30. doi:10.1007/s00115-020-01036-2
18. Alldredge BK, Gelb AM, Isaacs SM et al.: A Comparison of Lorazepam, Diazepam, and Placebo for the Treatment of Out-of-Hospital Status Epilepticus. *N Engl J Med* 2001; 345(9): 631–7. doi:10.1056/NEJMoa002141
19. Meyfroidt G, Baguley IJ, Menon DK: Paroxysmal sympathetic hyperactivity: the storm after acute brain injury. *Lancet Neurol* 2017; 16(9): 721–9. doi:10.1016/S1474-4422(17)30259-4
20. Rajajee V, Riggs B, Seder DB: Emergency Neurological Life Support: Airway, Ventilation, and Sedation. *Neurocrit Care* 2017; 27(s1): 4–28. doi:10.1007/s12028-017-0451-2
21. Goyal M, Menon BK, Van Zwam WH, et al.: Endovascular thrombectomy after large-vessel ischaemic stroke: A meta-analysis of individual patient data from five randomised trials. *Lancet* 2016; 387(10029): 1723–31. doi:10.1016/S0140-6736(16)00163-X
22. Jumaa MA, Zhang F, Ruiz-Ares G et al.: Comparison of Safety and Clinical and Radiographic Outcomes in Endovascular Acute Stroke Therapy for Proximal Middle Cerebral Artery Occlusion With Intubation and General Anesthesia Versus the Nonintubated State. *Stroke* 2010; 41(6): 1180–4. doi:10.1161/STROKEAHA.109.574194
23. Davis MJ, Menon BK, Baghirzada LB et al.: Anesthetic Management and Outcome in Patients during Endovascular Therapy for Acute Stroke. *Anesthesiology* 2012; 116(2): 396–405. doi:10.1097/ALN.0b013e318242a5d2
24. Berkhemer OA, van den Berg LA, Franzen PSS et al.: The effect of anesthetic management during intra-arterial therapy for acute stroke in MR CLEAN. *Neurology* 2016; 87(7): 656–64. doi:10.1212/WNL.0000000000002976
25. Abou-Chebl A, Yeatts SD, Yan B et al.: Impact of General Anesthesia on Safety and Outcomes in the Endovascular Arm of Interventional Management of Stroke (IMS) III Trial. *Stroke* 2015; 46(8): 2142–8. doi:10.1161/STROKEAHA.115.008761
26. Schönenberger S, Uhlmann L, Hacke W et al.: Effect of Conscious Sedation vs General Anesthesia on Early Neurological Improvement Among Patients With Ischemic Stroke Undergoing Endovascular Thrombectomy.

- JAMA 2016; 316(19): 1986. doi:10.1001/jama.2016.16623
27. Simonsen CZ, Yoo AJ, Sørensen LH et al.: Effect of General Anesthesia and Conscious Sedation During Endovascular Therapy on Infarct Growth and Clinical Outcomes in Acute Ischemic Stroke. *JAMA Neurol* 2018; 75(4): 470. doi:10.1001/jamaneuro.2017.4474
 28. Löwhagen Hendén P, Rentzos A, Karlsson J-E et al.: General Anesthesia Versus Conscious Sedation for Endovascular Treatment of Acute Ischemic Stroke. *Stroke* 2017; 48(6): 1601–7. doi:10.1161/STROKEAHA.117.016554
 29. Schönenberger S, Hendén PL, Simonsen CZ et al.: Association of General Anesthesia vs Procedural Sedation With Functional Outcome Among Patients With Acute Ischemic Stroke Undergoing Thrombectomy. *JAMA* 2019; 322(13): 1283. doi:10.1001/jama.2019.11455
 30. Ringleb PA et al.: S2e Leitlinie zur Akuttherapie des ischämischen Schlaganfalls. Leitlinien für Diagnostik und Ther der Neurol 2021
 31. Esmaeeli S, Valencia J, Buhl LK et al.: Anesthetic management of unruptured intracranial aneurysms: a qualitative systematic review. *Neurosurg Rev* 2021; 44(5): 2477–92. doi:10.1007/s10143-020-01441-w
 32. Fambrough DM, Drachman DB, Sathyamurti S: Neuromuscular Junction in Myasthenia Gravis: Decreased Acetylcholine Receptors. *Science* (80–) 1973; 182(4109): 293–5. doi:10.1126/science.182.4109.293
 33. Blichfeldt-Lauridsen L, Hansen BD: Anesthesia and myasthenia gravis. *Acta Anaesthesiol Scand* 2012; 56(1): 17–22. doi:10.1111/j.1399-6576.2011.02558.x
 34. Feldman JM: Cardiac Arrest after Succinylcholine Administration in a Pregnant Patient Recovered From Guillain-Barré Syndrome. *Anesthesiology* 1990; 72(5): 942–4. doi:10.1097/0000542-199005000-00026
 35. Kong KL, Willatts SM, Prys-Roberts C: Isoflurane compared with midazolam for sedation in the intensive care unit *BMJ* 1989; 298(6683): 1277–80. doi:10.1136/bmj.298.6683.1277
 36. Sackey PV, Martling C-R, Carlswärd C et al.: Short- and long-term follow-up of intensive care unit patients after sedation with isoflurane and midazolam – A pilot study*. *Crit Care Med* 2008; 36(3): 801–6. doi:10.1097/CCM.0B013E3181652FEE
 37. Bösel J, Purruicker JC, Nowak F et al.: Volatile isoflurane sedation in cerebrovascular intensive care patients using AnaConDa®: effects on cerebral oxygenation, circulation, and pressure. *Intensive Care Med* 2012; 38(12): 1955–64. doi:10.1007/s00134-012-2708-8
 38. Roberts DJ, Hall RI, Kramer AH et al.: Sedation for critically ill adults with severe traumatic brain injury: A systematic review of randomized controlled trials*. *Crit Care Med* 2011; 39(12): 2743–51. doi:10.1097/CCM.0b013e318228236f
 39. Rossetti AO, Milligan TA, Vulliëmoz S et al.: A randomized trial for the treatment of refractory status epilepticus. *Neurocrit Care* 2011; 14(1): 4–10. doi:10.1007/s12028-010-9445-z
 40. Claassen J, Hirsch LJ, Emerson RG, Mayer SA: Treatment of refractory status epilepticus with pentobarbital, propofol, or midazolam: A systematic review. *Epilepsia* 2002; 43(2): 146–53. doi:10.1046/j.1528-1157.2002.28501.x
 41. Prabhakar H, Kalaivani M: Propofol versus thiopental sodium for the treatment of refractory status epilepticus. *Cochrane Database Syst Rev* 2017; 2017(2). doi:10.1002/14651858.CD009202.pub4
 42. Thompson SA, Hantus S: Highly Epileptiform Bursts Are Associated With Seizure Recurrence. *J Clin Neurophysiol*. 2016; 33(1): 66–71. doi:10.1097/WNP.0000000000000232
 43. Prisco L, Ganau M, Aurangzeb S et al.: A pragmatic approach to intravenous anaesthetics and electroencephalographic endpoints for the treatment of refractory and super-refractory status epilepticus in critical care. *Seizure* 2020; 75: 153–64. doi:10.1016/j.seizure.2019.09.011
 44. Claassen J, Hirsch LJ, Emerson RG, Mayer SA: Treatment of Refractory Status Epilepticus with Pentobarbital, Propofol, or Midazolam: A Systematic Review. *Epilepsia* 2002; 43(2): 146–53. doi:10.1046/j.1528-1157.2002.28501.x
 45. Alvarez V, Lee JW, Westover MB et al.: Therapeutic coma for status epilepticus. *Neurology* 2016; 87(16): 1650–9. doi:10.1212/WNL.0000000000003224
 46. Sutter R, De Marchis GM, Semmlack S et al.: Anesthetics and Outcome in Status Epilepticus: A Matched Two-Center Cohort Study. *CNS Drugs* 2017; 31(1): 65–74. doi:10.1007/s40263-016-0389-5
 47. Marchi NA, Novy J, Faouzi M et al.: Status Epilepticus. *Crit Care Med* 2015; 43(5): 1003–9. doi:10.1097/CCM.0000000000000881
 48. Kowalski RG, Ziai WC, Rees RN et al.: Third-line antiepileptic therapy and outcome in status epilepticus. *Crit Care Med* 2012; 40(9): 2677–84. doi:10.1097/CCM.0b013e3182591ff1
 49. Sutter R, Semmlack S, Spiegel R et al.: Distinguishing in-hospital and out-of-hospital status epilepticus: clinical implications from a 10-year cohort study. *Eur J Neurol* 2017; 24(9): 1156–65. doi:10.1111/ene.13359
 50. Gaspard N, Foreman BP, Alvarez V et al.: New-onset refractory status epilepticus. *Neurology* 2015; 85(18): 1604–13. doi:10.1212/WNL.0000000000001940
 51. Meierkord H, Boon P, Engelsens B et al.: EFNS guideline on the management of status epilepticus in adults. *Eur J Neurol* 2010; 17(3): 348–55. doi:10.1111/j.1468-1331.2009.02917.x
 52. Rossetti AO, Logroscino G, Bromfield EB: Refractory status epilepticus: effect of treatment aggressiveness on prognosis. *Arch Neurol* 2005; 62(11): 1698–702. doi:10.1001/archneur.62.11.1698
 53. Opdenakker O, Vanstraelen A, De Sloovere V, Meyfroidt G: Sedatives in neurocritical care: an update on pharmacological agents and modes of sedation. *Curr Opin Crit Care* 2019; 25(2): 97–104. doi:10.1097/MCC.0000000000000592
 54. LeBlanc JM, Dasta JF, Kane-Gill SL: Role of the Bispectral Index in Sedation Monitoring in the ICU. *Ann Pharmacother* 2006; 40(3): 490–500. doi:10.1345/aph.1E491
 55. Marklund N: The neurological wake-up test—a role in neurocritical care monitoring of traumatic brain injury Patients? *Front Neurol* 2017; 8(OCT): 1–8. doi:10.3389/fneur.2017.00540
 56. Helbok R, Kurtz P, Schmidt MJ et al.: Effects of the neurological wake-up test on clinical examination, intracranial pressure, brain metabolism and brain tissue oxygenation in severely brain-injured patients. *Crit Care* 2012; 16(6): R226. doi:10.1186/cc11880
 57. Esnault P, Montcriol A, D’Aranda E et al.: Early neurological wake-up test in intubated brain-injured patients: A long-term, single-centre experience. *Aust Crit Care* 2017; 30(5): 273–8. doi:10.1016/j.aucc.2016.10.002
 58. Skoglund K, Enblad P, Marklund N: Effects of the neurological wake-up test on intracranial pressure and cerebral perfusion pressure in brain-injured patients. *Neurocrit Care* 2009; 11(2): 135–42. doi:10.1007/s12028-009-9255-3
 59. Skoglund K, Hillered L, Purins K et al.: The neurological wake-up test

does not alter cerebral energy metabolism and oxygenation in patients with severe traumatic brain injury. *Neurocrit Care* 2014; 20(3): 413–26. doi:10.1007/s12028–013–9876–4

60. Skoglund K, Enblad P, Hillered L, Marklund N: The neurological wake-up test increases stress hormone levels in patients with severe traumatic brain injury. *Crit Care Med* 2012;

40(1): 216–22.
doi:10.1097/CCM.0b013e31822d7dbdxx